

Freeze-out parameters from electric charge and baryon number fluctuations: is there consistency?

S. Borsanyi¹, Z. Fodor^{1,2,3}, S. D. Katz^{2,4}, S. Krieg^{1,3}, C. Ratti⁵, K. K. Szabo^{1,3}

¹ *Department of Physics, Wuppertal University,
Gaussstr. 20, D-42119 Wuppertal, Germany*

² *Inst. for Theoretical Physics, Eötvös University,
Pázmány P. sétány 1/A, H-1117 Budapest, Hungary*

³ *Jülich Supercomputing Centre,
Forschungszentrum Jülich, D-52425 Jülich, Germany*

⁴ *MTA-ELTE "Lendület" Lattice Gauge Theory Research Group,
Pázmány P. sétány 1/A, H-1117 Budapest, Hungary*

⁵ *Dip. di Fisica, Università di Torino and INFN, Sezione di Torino
via Giuria 1, I-10125 Torino, Italy*

(Dated: March 26, 2014)

Recent results for moments of multiplicity distributions of net-protons and net-electric charge from the STAR collaboration are compared to lattice QCD results for higher order fluctuations of baryon number and electric charge by the Wuppertal-Budapest collaboration, with the purpose of extracting the freeze-out temperature and chemical potential. All lattice simulations are performed for a system of 2+1 dynamical quark flavors, at the physical mass for light and strange quarks; all results are continuum extrapolated. We show that it is possible to extract an upper value for the freeze-out temperature, as well as precise baryo-chemical potential values corresponding to the four highest collision energies of the experimental beam energy scan. Consistency between the freeze-out parameters obtained from baryon number and electric charge fluctuations is found. The freeze-out chemical potentials are now in agreement with the statistical hadronization model.

The fundamental theory of strong interactions predicts a transition from a hadronic system to a partonic one at sufficiently high temperatures or densities; this transition is an analytic crossover, as shown by lattice QCD simulations [1]. The conditions of temperature or density needed to create the deconfined phase of QCD can be reached in the laboratory: relativistic heavy ion collisions are contributing tremendously to our understanding of the QCD phase diagram and the properties of the quark-gluon plasma (QGP). The precision reached in the most recent lattice QCD simulations, as well as the increasing availability of data from the heavy ion experimental programs, allow us to compare the theoretical and experimental results in a very efficient way.

Fluctuations of conserved charges (electric charge, baryon number and strangeness) are certainly a major example of this fruitful synergy. These observables, especially the higher order moments, were originally proposed as a signature for the QCD critical point [2–4], namely the point of the phase diagram which marks the change from crossover to first-order phase transition. As a consequence, experimental results for moments of net-electric charge and net-proton multiplicity distributions have recently been published, in the collision-energy range $\sqrt{s} = 7.7\text{--}200$ GeV covered by the RHIC beam energy scan [5–8]. The recent idea of extracting the freeze-out parameters of a heavy-ion collision through a comparison between lattice QCD results and experimental data has renewed the interest towards these observables even at $\mu = 0$ [9–11].

The chemical freeze-out, defined as the last inelastic scattering of hadrons before detection, has already been studied in terms of the statistical hadronization model by fitting a chemical potential and a temperature parameter to the pion, kaon, proton and other accessible yields from experiment [12–15]. By decreasing the collision energy, the freeze-out chemical potential increases; repeating the analysis for a series of beam energies provides a freeze-out curve in the (T, μ) plane.

The comparison between the experimental and lattice QCD results for the electric charge and baryon number fluctuations allows a first-principle determination of the freeze-out temperature and chemical potential, under the assumption that the experimentally measured fluctuations can be described in terms of the equilibrium system simulated on the lattice. Possible experimental sources of non-thermal fluctuations are corrected for in the STAR data analysis: the centrality-bin-width correction method minimizes effects due to volume variation because of finite centrality bin width; the moments are corrected for the finite reconstruction efficiency based on binomial probability distribution [16]; the spallation protons are removed by appropriate cuts in p_T [8]. In general, effects due to kinematic cuts are found to be small [17]. Besides, final-state interactions in the hadronic phase and non-equilibrium effects might become relevant and affect fluctuations [18, 19], therefore a fundamental check in favor of the equilibrium scenario is the consistency between the freeze-out parameters yielded by different quantum numbers e.g. electric charge and baryon

number. In particular, while the freeze-out temperature might be flavor-dependent [20], the chemical potentials as a function of the collision energy should be the same for all species. The present level of precision reached by lattice QCD results, obtained at physical quark masses and continuum-extrapolated, allows to perform this check for the first time.

One more caveat is in order, since experimentally only the net-proton multiplicity distribution is measured, as opposed to the lattice net-baryon number fluctuations. Recently it was shown that, once the effects of resonance feed-down and isospin randomization are taken into account [21, 22], the net-proton and net-baryon number fluctuations are numerically very similar [23].

In this paper we show for the first time that it is possible to find a consistency between the freeze-out parameters yielded by electric charge and baryon number fluctuations. This is achieved by systematically comparing our continuum-extrapolated results for higher order fluctuations of these conserved charges [11] to the corresponding experimental data by the STAR collaboration at RHIC [7, 8]. We are using the newly published, efficiency-corrected experimental results for the net-charge fluctuations and combine them with our lattice results presented in Ref. [11]. We also extract independent freeze-out conditions from the net-proton fluctuations and systematically compare the outcomes of the two. Details of the lattice simulations can be found in [11].

The fluctuations of baryon number, electric charge and strangeness are defined as

$$\chi_{lmn}^{BSQ} = \frac{\partial^{l+m+n}(p/T^4)}{\partial(\mu_B/T)^l \partial(\mu_S/T)^m \partial(\mu_Q/T)^n}; \quad (1)$$

they are related to the moments of the multiplicity distributions of the corresponding conserved charges. It is convenient to introduce the following, volume-independent ratios

$$\begin{aligned} \chi_3/\chi_2 &= S\sigma & \chi_4/\chi_2 &= \kappa\sigma^2 \\ \chi_1/\chi_2 &= M/\sigma^2 & \chi_3/\chi_1 &= S\sigma^3/M. \end{aligned} \quad (2)$$

The chemical potentials μ_B , μ_Q and μ_S are related in order to match the experimental situation: the finite baryon density in the system is due to light quarks only, since it is generated by the nucleon stopping in the collision region. The strangeness density $\langle n_S \rangle$ is then equal to zero for all collision energies, as a consequence of strangeness conservation. Besides, the electric charge and baryon-number densities are related, in order to match the isospin asymmetry of the colliding nuclei: $\langle n_Q \rangle = Z/A \langle n_B \rangle$. $Z/A = 0.4$ represents a good approximation for Pb-Pb and Au-Au collisions.

As a consequence, μ_Q and μ_S depend on μ_B so that these conditions are satisfied. This is achieved by Taylor-expanding the densities in these three chemical potentials

up to μ_B^3 [10]:

$$\begin{aligned} \mu_Q(T, \mu_B) &= q_1(T)\mu_B + q_3(T)\mu_B^3 + \dots \\ \mu_S(T, \mu_B) &= s_1(T)\mu_B + s_3(T)\mu_B^3 + \dots \end{aligned} \quad (3)$$

Our continuum extrapolated results for the functions $q_1(T)$, $q_3(T)$, $s_1(T)$, $s_3(T)$ were shown in [11]. The quantities that we consider to extract the freeze-out T and μ_B , are the ratios $R_{31}^B = \chi_3^B/\chi_1^B$ and $R_{12}^B = \chi_1^B/\chi_2^B$ respectively, at values of (μ_B, μ_Q, μ_S) , which satisfy the physical conditions discussed in the previous paragraph. As shown in Ref. [11], the leading order in χ_3^B/χ_1^B is independent of μ_B , while the LO in χ_2^B/χ_1^B is linear in μ_B . This allows us to use R_{31}^B to extract the freeze-out temperature; the ratio R_{12}^B is then used to extract μ_B (notice that our results for R_{12}^B are obtained up to NLO in μ_B).

We then independently extract μ_B from χ_1^Q/χ_2^Q (which is also linear in μ_B to LO), in order to check whether different conserved charges yield consistent freeze-out parameters. In Ref. [11], we compared the lattice results for χ_3^Q/χ_1^Q to the preliminary, efficiency-uncorrected data from the STAR collaboration, to extract an upper limit for the freeze-out temperature. We then obtained the corresponding chemical potentials by performing the same kind of comparison for χ_1^Q/χ_2^Q . The new, efficiency-corrected results for the moments of the net-charge multiplicity distribution from STAR show significant differences, compared to the uncorrected ones. This yields different values for μ_B , compared to the ones obtained in [11]. As for χ_3^Q/χ_1^Q , the experimental uncertainty on the corrected data is such that presently it is not possible




FIG. 1. R_{31}^B : the colored symbols show finite- N_t lattice QCD results. The continuum extrapolation is represented by black points (from Ref. [11]). The dark-orange band shows the recent experimental measurement by the STAR collaboration [7]: it was obtained by averaging the 0-5% and 5-10% data at the four highest energies ($\sqrt{s} = 27, 39, 62.4, 200$ GeV). The green-shaded area shows the valid temperature range.

to extract a meaningful freeze-out temperature from this observable.

In Fig. 1 we show the comparison between the lattice results for $\frac{\chi_3^B(T, \mu_B)}{\chi_1^B(T, \mu_B)}$ and the experimental measurement of $S_p \sigma_p^3 / M_p$ by the STAR collaboration [7]. The latter has been obtained for a 0-10% centrality, at the four highest energies ($\sqrt{s} = 27, 39, 62.4, 200$ GeV). Since the curvature of the phase diagram is small around $\mu_B = 0$ [24], this average allows to determine the freeze-out temperature. The green-shaded area shows the valid temperature range: due to the uncertainty on the lattice results in the low-temperature regime, it is only possible to extract an upper value for the freeze-out temperature: the freeze-out takes place at a temperature $T_f \lesssim 148$ MeV, which is somewhat lower than expected from previous analyses [25] (allowing for a two-sigma deviation for the lattice simulations and the experimental measurements, the highest possible T_f is 151 MeV). In Refs. [26, 27] we have published the lattice determination of the transition temperature from various chiral observables in the range 147-157 MeV. For the minimum of the speed of sound we found 145(5) MeV in [28]. The discussed freeze-out temperature is thus in the cross-over region around or slightly below the central value.

We now proceed to determine the freeze-out chemical potential μ_B , by comparing the lattice results for R_{12}^B and R_{12}^Q (as functions of the chemical potential, and in the temperature range ($140 \leq T_f \leq 150$) MeV) to the experimental results for M_p/σ_p^2 and M_Q/σ_Q^2 published by the STAR collaboration in Refs. [7, 8, 29]. This comparison is shown in the two panels of Fig. 2: the two quantities allow for an independent determination of μ_B from electric charge and baryon number: the corresponding values are listed in Table I, and shown in Fig. 3. Consistency between the two values of baryon-chemical potential is found for all collision energies (the non-monotonicity of the lattice results for R_{12}^B at $\mu_B \geq 130$ MeV does not allow a determination of μ_B from this observable at $\sqrt{s} = 27$ GeV). Let us now compare the chemical potentials in Table I to those found earlier in statistical fits [12, 14, 30]. Plotting the parametrization of Refs. [12, 30] together with our values we find a remarkable agreement (see Fig. 3). Note that, for the freeze-out temperature, statistical models typically yield a somewhat higher value: e.g. 164 MeV in Refs. [12, 31]. Towards the lower end in temperature range we find Ref. [32] with $T_f = 155 \pm 8$ MeV with $\mu_B = 25 \pm 1$ MeV, at $\sqrt{s} = 200$ GeV.

The comparison of our lattice results to the latest efficiency-corrected STAR data hints at a consistency of the freeze-out chemical potential if we assume an agreement in the temperature. This assumption was well motivated by the proton and charge skewness data. Let us now take the assumption further: if the freeze-out can be described by the same temperature and chemical poten-




FIG. 2. Upper panel: R_{12}^B as a function of μ_B . The three points correspond to the STAR data for M_p/σ_p^2 at collision energies $\sqrt{s} = 39, 62.4, 200$ GeV and centrality 0-10%, from Ref. [7] (the $\sqrt{s} = 27$ GeV point is also shown, but the non-monotonicity of the lattice results at $\mu_B \geq 130$ MeV does not allow a determination of μ_B from it). Lower panel: R_{12}^Q as a function of μ_B . The four points correspond to the STAR data for M_Q/σ_Q^2 at $\sqrt{s} = 27, 39, 62.4, 200$ GeV and centrality 0-10%, from Ref. [8]. In both panels, the colored symbols correspond to the lattice QCD results in the continuum limit, for the range ($140 \leq T_f \leq 150$) MeV. The arrows show the extracted values for μ_B at freeze-out.

tials for charge and protons, then one can create a combined observable: $R_{12}^Q/R_{12}^B = [M_Q/\sigma_Q^2]/[M_B/\sigma_B^2]$. Here, the volume factor of the charge and baryon (proton) measurements cancel separately. Should our assumption be correct, this ratio of ratios is the preferable thermometer: it is far easier to obtain both for lattice and experiment since it does not involve skewness or kurtosis. We have lattice data available to $\sim \mu_B^2$ order, which we use when comparing our results to data. Such a comparison is shown in Fig. 4. Contrary to the skewness thermometer, here we see a clear monotonic temperature dependence without the hardly controllable lattice errors at low temperatures. This allows for the identification of a narrow

\sqrt{s} [GeV]	μ_B^f [MeV] (from B)	μ_B^f [MeV] (from Q)
200	25.6 ± 2.4	22.6 ± 2.4
62.4	69 ± 5.7	65.9 ± 7.2
39	104 ± 10	100 ± 9
27	-	134.5 ± 12.5

TABLE I. Freeze-out μ_B vs. \sqrt{s} , for the four highest-energy STAR measurements. The μ_B values and error-bars in this table have been obtained under the assumption that $140 \text{ MeV} \leq T_f \leq 150 \text{ MeV}$. This uncertainty dominates the overall errors. Other (minor) sources of uncertainty are the lattice statistics and the experimental error.




FIG. 3. Freeze-out chemical potential μ_B as a function of the collision energy. The red stars show the μ_B obtained by fitting R_{12}^Q , the blue squares have been obtained by fitting R_{12}^B . The black curve comes from the Statistical Hadronization Model analysis of Refs. [12, 30].

temperature band, instead of an upper limit.

The thermometer in Fig. 4 is, in fact, a consistency criterion: it agrees to the experimental data at the temperature which needs to be assumed for the freeze-out if the proton and charge fluctuations reflect the grand canonical ensemble at a common chemical potential. For high enough energies ($\sqrt{s} \geq 39 \text{ GeV}$) this consistency is granted if freeze-out occurs in the range $T_f = 144 \pm 6 \text{ MeV}$. Notice that this temperature range lies just below the upper limit that we determined independently in Fig. 1.

In conclusion, in the present paper we have extracted the freeze-out conditions (temperature and chemical potential) by comparing our continuum extrapolated lattice QCD results to the experimental moments of net-charge and net-proton multiplicity distribution by the STAR collaboration. These new, efficiency corrected, experimental data point at a lower freeze-out temperature compared to previous estimates. This is compatible with the expectation that the freeze-out should occur just below the transition [33]. The independent determinations




FIG. 4. R_{12}^Q/R_{12}^B : the colored symbols correspond to continuum-extrapolated lattice QCD simulations at different values of μ_B . The dashed lines show the recent experimental measurements by the STAR collaboration [7, 8] for a 0-10% centrality and different collision energies. The green-shaded area shows the valid temperature range, $T_f = (144 \pm 6) \text{ MeV}$.

of the freeze-out chemical potentials from electric charge and baryon number show a remarkable consistency with each other. This comparison is possible for the first time, and the consistency of the results is of fundamental importance to validate the hypothesis on which this method is based, namely that the experimentally created system is close to thermal equilibrium at the freeze-out and can be described by lattice QCD simulations, at least in the light quark sector.

Acknowledgments: C. Ratti acknowledges fruitful discussions with Francesco Becattini, Rene Bellwied and Bill Llope. This project was funded by the DFG grant SFB/TR55. The work of C. Ratti is supported by funds provided by the Italian Ministry of Education, Universities and Research under the Fibr Research Grant RBFR0814TT. S. D. Katz is funded by the ERC grant ((FP7/2007-2013)/ERC No 208740) as well as the "Lendület" program of the Hungarian Academy of Sciences ((LP2012-44/2012). The numerical simulations were performed on the QPACE machine, the GPU cluster at the Wuppertal University and on JUQUEEN (the Blue Gene/Q system of the Forschungszentrum Juelich).

-
- [1] Y. Aoki, G. Endrodi, Z. Fodor, S. D. Katz and K. K. Szabo, *Nature* **443**, 675 (2006) [hep-lat/0611014].
 - [2] M. A. Stephanov, K. Rajagopal and E. V. Shuryak, *Phys. Rev. D* **60**, 114028 (1999) [hep-ph/9903292].
 - [3] R. V. Gavai and S. Gupta, *Phys. Rev. D* **78**, 114503 (2008) [arXiv:0806.2233 [hep-lat]].
 - [4] M. Cheng, *et al.*, *Phys. Rev. D* **77**, 014511 (2008)

- [arXiv:0710.0354 [hep-lat]].
- [5] D. McDonald [STAR Coll.], arXiv:1210.7023 [nucl-ex].
- [6] N. R. Sahoo [STAR Coll.], arXiv:1212.3892 [nucl-ex].
- [7] L. Adamczyk *et al.* [STAR Collaboration], arXiv:1309.5681 [nucl-ex].
- [8] L. Adamczyk *et al.* [STAR Collaboration], arXiv:1402.1558 [nucl-ex].
- [9] F. Karsch, Central Eur. J. Phys. **10**, 1234 (2012) [arXiv:1202.4173 [hep-lat]].
- [10] A. Bazavov, *et al.*, Phys. Rev. Lett. **109**, 192302 (2012) [arXiv:1208.1220 [hep-lat]].
- [11] S. Borsanyi, Z. Fodor, S. D. Katz, S. Krieg, C. Ratti and K. K. Szabo, Phys. Rev. Lett. **111** (2013) 062005 [arXiv:1305.5161 [hep-lat]].
- [12] A. Andronic, P. Braun-Munzinger and J. Stachel, Phys. Lett. B **673** (2009) 142 [Erratum-ibid. B **678** (2009) 516] [arXiv:0812.1186 [nucl-th]].
- [13] F. Becattini, J. Manninen and M. Gazdzicki, Phys. Rev. C **73**, 044905 (2006)
- [14] J. Cleymans, H. Oeschler, K. Redlich and S. Wheaton, Phys. Rev. C **73** (2006) 034905 [hep-ph/0511094].
- [15] J. Manninen and F. Becattini, Phys. Rev. C **78**, 054901 (2008) [arXiv:0806.4100 [nucl-th]].
- [16] A. Bzdak and V. Koch, Phys. Rev. C **86** (2012) 044904 [arXiv:1206.4286 [nucl-th]].
- [17] P. Alba, W. Alberico, R. Bellwied, M. Bluhm, V. M. Sarti, M. Nahrgang and C. Ratti, arXiv:1403.4903 [hep-ph].
- [18] J. Steinheimer, J. Aichelin and M. Bleicher, Phys. Rev. Lett. **110** (2013) 042501 [arXiv:1203.5302 [nucl-th]].
- [19] F. Becattini, M. Bleicher, T. Kollegger, T. Schuster, J. Steinheimer and R. Stock, Phys. Rev. Lett. **111** (2013) 082302 [arXiv:1212.2431 [nucl-th]].
- [20] R. Bellwied, S. Borsanyi, Z. Fodor, S. DKatz and C. Ratti, Phys. Rev. Lett. **111** (2013) 202302 [arXiv:1305.6297 [hep-lat]].
- [21] M. Kitazawa and M. Asakawa, Phys. Rev. C **85** (2012) 021901 [arXiv:1107.2755 [nucl-th]].
- [22] M. Kitazawa and M. Asakawa, Phys. Rev. C **86** (2012) 024904 [Erratum-ibid. C **86** (2012) 069902] [arXiv:1205.3292 [nucl-th]].
- [23] M. Nahrgang, M. Bluhm, P. Alba, R. Bellwied and C. Ratti, arXiv:1402.1238 [hep-ph].
- [24] G. Endrodi, Z. Fodor, S. D. Katz and K. K. Szabo, JHEP **1104**, 001 (2011) [arXiv:1102.1356 [hep-lat]].
- [25] S. Mukherjee and M. Wagner, PoS CPOD **2013** (2013) 039 [arXiv:1307.6255 [nucl-th]].
- [26] Y. Aoki, S. Borsanyi, S. Durr, Z. Fodor, S. D. Katz, S. Krieg and K. K. Szabo, JHEP **0906** (2009) 088 [arXiv:0903.4155 [hep-lat]].
- [27] S. Borsanyi *et al.* [Wuppertal-Budapest Collaboration], JHEP **1009** (2010) 073 [arXiv:1005.3508 [hep-lat]].
- [28] S. Borsanyi, G. Endrodi, Z. Fodor, A. Jakovac, S. D. Katz, S. Krieg, C. Ratti and K. K. Szabo, JHEP **1011** (2010) 077 [arXiv:1007.2580 [hep-lat]].
- [29] The efficiency-corrected data for the lowest cumulant ratio (c_2/c_1) for net-protons can be found on the public STAR webpage.
- [30] A. Andronic, P. Braun-Munzinger and J. Stachel, Nucl. Phys. A **772** (2006) 167 [nucl-th/0511071].
- [31] J. Adams *et al.* [STAR Collaboration], Nucl. Phys. A **757** (2005) 102 [nucl-ex/0501009].
- [32] J. Rafelski, J. Letessier and G. Torrieri, Phys. Rev. C **72** (2005) 024905 [nucl-th/0412072].
- [33] P. Braun-Munzinger, J. Stachel and C. Wetterich, Phys. Lett. B **596** (2004) 61